

# Potato



## ***Healthier Plant, Higher Production***

Green Earth advisory works towards creating a Healthy Plant and a Good Soil structure. This acts as the basis for higher production, better survival under biotic & abiotic stress and lesser attack of plant and disease. We focus on improving the 3 core dimension:

- Soil Eco System.
- Plant Root System.
- Plant's structure & defence System.

We acknowledge the role of natural forces, but we firmly believe that this strategy easily handles over 80% challenges.

### ***Conductive Soil Eco System***

The recommended ready mix consortia of compatible biological agents specially blended with bio-enzymes and essential nutrients improves the soil eco-system in multiple modes.

- a) **Maximizes Nutrition Utilization** by solubilisation & mobilization of nutrients like Phosphorus, Potash, Zinc, Calcium, Magnesium, Sulphur, Iron, Manganese etc. Also improves Nitrogen availability through fixation from the atmosphere. Overall impact is on improvement of nutrient use efficiency.
- b) **Control of soil based pathogens** for better control of *Wilt, tuber Rot & Nematodes*. They cause maximum damage to the plant leading to lower production. Over 80% reductions in infestation are observed.
- c) **Improving the Soil Structure & Health** for better aeration and soil aggregation aiding the growth of beneficial microorganisms in soil.

**Recommendation:**

<b>Product</b>	<b>Dose /ac</b>	<b>Product</b>	<b>Dose /ac</b>
Reap NPK1	1000 ml	Sumona	250 – 500 ml
Reap ZFeMn	500 ml	Vima	250 – 500 ml

**Application: - With FYM:** Mix the products in 500-600 kg of FYM/Compost and apply on the field at the time bed preparation. **Drench:** Mix the products in 200 lit of water and drench on the field 20 days after planting.

***Nature of Soil Best Suited:***

A well drained, friable, porous with good organic content is ideal

***[NPK level in soil recommended: N 220-280 kg/ha, P 15-20 Kg/ha, K 180-280 Kg/ha]***

***Powerful Root Zone***

Proper absorption of nutrition from soil is possible only through a vigorous micro root network, robust and well spread root system with high surface area of the root.

R-Jaal which is recommendation develops up to 30 to 50% higher micro root structure. It creates a network (Jaal) of fine filaments that associate with plant roots and extend the roots far into the soil resources; much beyond the area of nutrient depletion. Thus it enables the plant to draw nutrients and water from the soil that a plant would not be able to access otherwise. Naturally, this helps plant survive in multiple stresses and make it strongly rooted in the soil.

Powerful root zone also helps in better development of tuber size, weight & quality.

**Recommendation: R-Jaal- 2 Kgs per Acre**

**Application:** Mix R-Jaal in 400-500 kg FYM/Compost and apply on the field at the time of bed preparation.

## **Robust Plant Structure**

Goal of commercial farming is highest production at optimum cost. Total photosynthesis capacity of the plant determines its production capacity. A well erected plant combined with good quality leaves only will have excellent photosynthesis capacity.

With this in mind we focus on delivering a robust plant structure and leaves quality,

- A. We recommend Bio-available Silica to create strong Cell Wall which leads to **stronger & bigger branches and stem**. It also increases the leaf size and makes it more erect.
- B. It also increases the uptake and transports more water, nutrients and plant secretions throughout the plant body. This availability of nutrition is critical for higher production.
- C. Plant Growth Regulator to increase the no of leaves, improvement in chlorophyll level for higher photosynthesis and other metabolic actions.

The above approach also builds a strong defence wall to block the pests from attacking the plant. This works as a powerful **defence against fungus, mites & sucking pest**. This is drastically reduces the cost & efforts incurred in pest management.

Stronger Plant structure also enables the plant to bear more fruits.

**Recommendations: SilPot + Green Flush** [4 to 5 sprays]

### **Application:**

Mix **SilPot** (2 ml / lit) + **Green Flush** (1- 1.5 ml / lit) and take spray. First spray 20 - 25 days after plantation. Repeat every 20-25 days interval up to rhizome development stage.

Spray **SilPot** (2 ml/lit) 1 to 2 days before using Weedicide.

## **Nutrition Management**

Nutrients (Fertilizers) are the food for the plant. It needs food for production. Even if all the factors are favourable, it still will require sufficient nutrition to support the final production.

A Green Earth practice optimizes and maximizes the utilization of available nutrition leading to higher production. In numbers it can be said to **improve nutrition utilization by 25-30%**

The table below explains the Recommended Doses of Fertilizer and Yield as per general farmer feedback. If all the above practices are implemented without changing the fertilizer doses, we have seen remarkable increase in yield. **Yield (Normal)** is with general practices as followed in the market, while **Yield (Green Earth)** is with above practice on Soil, Root and Plant Structure followed.

Component	Doses/ Acre	Yield (Normal)	Yield (Green Earth)
Nitrogen	60 Kgs	<b>6 - 7 MT</b>	<b>8 - 9 MT</b>
Phosphorous	30 Kgs		
Potash	50 Kgs		

The production can change due to the variety used, crop distance or farming under control environment (poly house / shed net etc). It is common sense that higher production will require higher amount of nutrition. We recommend you to follow nutrition chart provided by agronomy advisor.

#### **Stress Management – Biotic & Abiotic**

Biotic & Abiotic stress creates huge losses, sometime even wiping out the full crop.

The climate change has increased the Abiotic Stress to very high level. Early/Late Rain, intermittent dry spell, Heat Shock, Cold Shock, hail storm etc. With the above treatment, the plants can respond much better under such stress:

<b>Nature of Shock</b>	<b>Impact improvement</b>
Dry Spell Shock	Plants survive around <b>15 days</b> more, than controlled plot.
Heat Shock	Sustains up to <b>3°Celsius</b> & <b>10 days</b> more than controlled.
Cold Shock	Up to <b>10 days</b> more than controlled.
Very Heavy Rain	Will sustain longer and recover faster.
High Velocity Wind	<b>70%</b> lesser uprooting.
Hail Storm	<b>25%</b> lesser impact. Difficult to measure.
Weedicide Shock	<b>50%</b> lesser impact on Plant Health.

Abiotic stress also makes the plants weak and susceptible to pathogen attacks.

Biotic stress (commonly termed as Pest & Disease Attack) leads to losses from multiple angles, direct losses from the pest on health deterioration of the plant and Shock to the plant from the Pesticide use. Also the cost & labour incurred in spray of pesticides.

Green Earth recommended practices drastically reduce the chances of infection & infestation. The table illustrates the various impacts,

<b>Disease / Pest</b>	<b>% Risk Reduction in Infestation</b>
Tuber Rot, Nematode, Wilt	<b>80%</b> less chance of infection
Leaf Spot, Blight, Thrips & Mites	<b>50%</b> less chance of infection
Ring Spot, Tuber Moth	<b>25%</b> less chance of infection

In spite of all precautions, still if there is an attack we recommend taking advice from specialist and control accordingly.